

اثر تترودوتوکسین و کادمیم روی آزاد شدن و متابولیسم سیستم‌های

سروتونرژیک و نورآدرنژیک هیپوکامپ

عبدالوهاب وهاب زاده*

چکیده:

با استفاده از تکنیک میکرو دیالیز مغز، ارتباط بین تحریک نورونی و آزاد شدن و متابولیسم نوروترانسمیترهای هیپوکامپ مورد مطالعه قرار گرفت. تترودوتوکسین (TTX) بطور داخل هسته‌ای داده شد و یون کادمیم (Ca^{++}) با یون کلسیم (Ca^{++}) در CSF مصنوعی میکرودیالیز معاوضه گردید. اثرات این دو دارو روی سیستم‌های سروتونرژیک و نورآدرنژیک هیپوکامپ مطالعه شد.

موش سفید (۲۵۰ تا ۳۰۰ گرم) انتخاب و پراب‌های میکرودیالیز در هیپوکامپ قدامی در شرایط بیهوشی (کلیال هیدرات) کاسته شد. بعد از دوران رکاوری ۵- هیدروکسی تریپتامین (5-HT) ۵- هیدارکسی اندول استیک اسید (5-HIAA) نورآدرنالین (NA) و دی هیدراکسی فنیل اسید استیک (DOPAC) با استفاده از HPLC با آشکارساز الکتروشیمیایی اندازه‌گیری شد. از مسدودکننده‌های برداشت HT 5- (Citalopram) و (Dssipramine) NA نیز استفاده شد.

بعد از ثبوت خط پایه و برداشت سه نمونه شاهد TTX داده شد و این سبب کاهش خط پایه 5-HT به $4(p < 0.05) \pm 29\%$ 5-HIAA به $7(p < 0.05) \pm 87\%$ NA به $3(p < 0.05) \pm 24\%$ و DOPAC به $17(p < 0.05) \pm 55\%$ در عرض ۱۰۰ دقیقه گردید ($n=6$) که جایگزین Ca^{++} گردید کاهش خط پایه 5-HT به $8(p < 0.05) \pm 38\%$ در عرض ۱۰۰ دقیقه گردید. ($n=6$).

نتایج نشان می‌دهد که آزاد شدن 5-HT و NA در هیپوکامپ روندی فعال است و بستگی به دشارژ شدن پایانه عصبی این سیستم‌ها و ورود یون Ca^{++} به این پایانه‌ها دارد. متابولیسم این سیستم‌ها متعاقب به آزاد شدن، در هر مورد کاهش می‌یابد. با این حال نتایج نیاز مبرم یون Ca^{++} برای فعال تر شدن آنزیم‌های متابولیکی این سیستم‌ها را توصیه می‌کند.

۳- کادمیم

۲- تترودوکسین

کلید واژه‌ها: ۱- میکرو دیالیز

۴- نورآدرنژیک هیپوکامپی ۵- سیستم‌های سروتونرژیک

* بخش فیزیولوژی، دانشکده پزشکی، دانشگاه علوم پزشکی و خدمات بهداشتی درمانی، ایران - تهران

THE EFFECT OF TETRODOTOXIN AND CADMIUM ON HIPPOCAMPAL NORADRENALINE AND 5-HYDROXYTRYPTAMINE RELEASE AND TURNOVER.

A. Vahabzadeh *

ABSTRACT

Using the technique of in vivo microdialysis, we examined the relationship between neuronal firing and neurotransmitter release and metabolism in the hippocampus. The neurotoxin tetrodotoxin (TTX) was applied locally to block voltage-sensitive sodium channels, and extracellular Ca^{++} was replaced with cadmium (Cd^{++}) to inhibit Ca^{++} dependent neurotransmitter release. The effects of these drugs on both noradrenergic and serotonergic transmitter system were evaluated. The results show that 5-HT and NA release in hippocampal nerve terminal are active process and depend on the cell firing and influx of Ca^{++} ions prior to release. 5-HT and NA synthesis and turnover are also decreased by TTX and Cd^{++} , these decays are greater in the absence of Ca^{++} ions. This suggests an essential role of Ca^{++} ions for hippocampal monoamine synthesis and turnover.

Key Words: 1) Microdialysis 2) tetrodotoxin 3) Cadmium
4) Hippocampal noradrenergic 5) serotonergic systems.

INTRODUCTION

The classical monoamines neurotransmitters 5-HT and NA are probably the most studied of all CNS transmitter substances. In the central nervous system both NA and

5-HT have been implicated in a wide range of behavioural and psychological functions such as aggressive and predatory behaviour, anxiety and depression. In spite of such

* Department of physiology, Faculty of medicine, Iran University of Medical Science, Tehran, Iran.

diverse functions attributed to 5-HT and NA, obtaining direct evidence has been hampered by the difficulty inherent in the *in vivo* estimation of these neurotransmitters. Electrophysiological recording from single units of both these systems in freely moving animals^(1, 32) has been utilised to evaluate the activity of these systems; this technique is unable to detect changes in the amount of transmitter release, synthesis, turnover resulting from changes in synaptic terminals. Various neurochemical approaches, such as push-pull perfusion^(12,30) and *in vivo* voltammetry^(3,22) have also been used to monitor the release of brain neurotransmitters. However, these techniques either produce tissue damage in the implanted area of brain^(13,25,30) because of the continuous tissue wash and the size of push-pull cannulae, or are unable to monitor extracellular NA and 5-HT levels directly, partly because of the similarity of the oxidation potential of the metabolites and the neurotransmitter^(3,4,19,21,31). Recently, the development of *in vivo* microdialysis techniques for direct measurement of the extracellular concentrations of the neurotransmitters and their metabolites have presented promising tools for investigating the conditions under which the activity of these neurotransmitter systems is altered as well as how this activity is regulated. The use of the dialysis technique also provided several advantages over other

in vivo techniques. The most important are its high resolution and the ability to measure several different analytes in the same sample in freely moving animal so that behaviour can be correlated with chemical changes in the brain. Instead of such advantages, brain dialysis is an invasive in nature. It attempts to monitor the release of transmitter from nerve terminals by inserting into the brain a probe several orders of magnitude larger than the biological structure than the biological structure under study. Thus due to this nature it is necessary to establish specific criteria for evaluation of neurotransmitter output as estimated by dialysis. These include the use of the selective voltage dependent Na⁺ channel blocker tetrodotoxin (TTX), replacement of Cd⁺⁺ with Ca⁺⁺ in perfusing medium for assessment of the physiological release process and activity of the synthetic enzyme^(9,16,29,33,34). The results obtained from these brain dialysis studies depends on at least three variables: type of probe, post implantation interval, and whether anaesthetised or freely moving animals are used.

MATERIAL & METHOD

Rats (250-300g) were implanted with microdialysis probes in the ventral hippocampus under chloral hydrate anaesthesia (500 mg/kg *i.p.*), and allowed to recover overnight. Samples of dialysate

were collected every 20 minutes from the unanaesthetised and freely moving animals and assayed for 5-HT, 5-HIAA, NA and DOPAC using HPLC with electrochemical detection. Two separate HPLC assay systems were used one of which measured 5-HT and 5-HIAA, and the other measured NA and DOPAC. Since basal levels of the transmitters were close to the limit of detection of the assay system, citalopram (μM), or desipramine ($10\ \mu\text{M}$) was added in most experiments to the perfusion medium for the assay of 5-HT and NA respectively; these drugs are selective uptake blockers for these transmitters. The neuronal origin of the compounds was verified either by replacement of Ca^{++} with Cd^{++} in the perfusing medium or by the use of the Na^{++} channel blocker TTX ($1\ \mu\text{M}$). The mean of the three basal samples was calculated and the value of all individual samples was expressed as a percentage of this mean. For statistical significance of the effect of a drug comparisons were made with the last sample before the addition of the drug using absolute values and the paired student's test.

RESULTS

Serotonergic nerve terminals

In order to determine to what extent the basal levels of 5-HT and 5-HIAA in hippocampal dialysate are impulse traffic-dependent the voltage-dependent

Na^{+} channel blocker TTX was infused. Local infusion of TTX ($1\ \mu\text{M}$) through the dialysis probe caused an initial rapid decrease in the concentration of 5-HT which was followed by a further slower reduction to $24\pm 5\%$ ($p<0.006$, $n=3$) by 80 min, the duration of experiment. The initial rapid reduction of 5-HT to $33\pm 6\%$ occurred in the first 20 min sample.

Local administration of TTX caused a gradual and much smaller reduction in the concentration of 5-HIAA to $84\pm 3\%$ ($p<0.001$, $n=8$) by 80 min (Fig.1.).

In order to determine the Ca^{++} -dependence of basal 5-HT and 5-HIAA, Ca^{++} was replaced with Cd^{++} in the Ringer solution. This caused an initial rapid decrease to $53\pm 18\%$ in the first 20 min sample in the concentration of 5-HT, was followed by a slower reduction to $23\pm 8\%$ ($p<0.01$, $n=3$) by 100 min; 5-HIAA was reduced to $18\pm 5\%$ ($p<0.003$, $n=6$) by 100 min (Fig.2.).

Noradrenergic nerve terminals

Local infusion of TTX ($1\ \mu\text{M}$) through the dialysis probe caused an initial rapid decrease to $49\pm 5\%$ within the first 20 min sample in the basal concentration of NA followed by a further slow reduction to $24\pm 3\%$ ($p<.02$, $n=3$) by 100 min. The concentration of DOPAC was reduced to $55\pm 17\%$ ($p<0.001$, $n=8$) by 100 min (Fig.3.).




Figure 1. The effect of TTX on the basal level of 5-HT and 5-HIAA. Changes in the concentration of 5-HT are shown as a percentage of the last sample taken before the infusion. The solid bar shows the infusion of TTX. * $p < 0.005$ ($n = 8$ for 5-HIAA and 3 for 5-HT) compared to the last sample taken before TTX.

Replacement of Ca^{++} with Cd^{++} in the perfusion medium caused an initial rapid decrease the basal concentration of NA followed by a progressive slow reduction to

$32 \pm 8\%$ ($p < 0.01$, $n = 3$) by 100 min. There was a more gradual reduction in DOPAC level to $47 \pm 5\%$ ($p < 0.0003$, $n = 6$) by 100 min (Fig.4).




Figure 2. The effect of the replacement of Ca^{++} with Cd^{++} in the perfusion medium on the basal level of 5-HT and d5-HIAA. The solid bar shows the infusion of Cd^{++} . * $p < 0.05$ ($n = 6$ for 5HIAA and 3 for 5-HT) compared to the last sample taken before Cd^{++} .




Figure 3. The effect of TTX on the basal level of NA and DOPAC. The solid bar shows the infusion of TTX. * $p < 0.05$ ($n = 8$ for DOPAC and 3 for NA) compared to the last sample taken before TTX.




Figure 4. The effect of the replacement of Ca^{++} with Cd^{++} in the perfusion medium on the basal level of NA and DOPAC. The solid bar shows the infusion of Cd^{++} . * $p < 0.05$ ($n=6$ for DOPAC and 3 for NA) compared to the last sample taken before Cd^{++}

DISCUSSION

In the present study in vivo microdialysis has been used to monitor the basal levels of hippocampal 5HT and NA their metabolites in unanaesthetised freely moving rats. The voltage-dependent Na^+ channel blocker, TTX, was used locally to determine to what extent the basal levels of 5-HT and NA are impulse-dependent. For the metabolites

TTX was used to determine the link between impulse traffic, transmitter synthesis and turnover. Previous studies have shown that TTX reduces the spontaneous release in vivo of acetylcholine (ACh) (6,7), DA (34), NA (33) and 5-HT (2,9,17,29), as well as the electrically evoked release of several amino acids, including GABA (15). The reduction to 24%

Downloaded from http://journals.tums.ac.ir/ at 11:44 IRDT on Sunday July 15th 2012

[Downloaded from rjms.tums.ac.ir on 2024-07-22]

of both 5-HT and NA by TTX confirms the dependence of their release on nerve impulse traffic as also found by others^(2,17,29). The remaining 5-HT and NA is probably due to the fact that TTX did not reach all the nerve terminals.

The smaller reduction of the two metabolites levels suggests that synthesis is less closely coupled to impulse traffic than release. There is however a considerable difference between metabolites, 5-HIAA showing a much smaller reduction after TTX than DOPAC. This suggests that in the NA pathway there is close coupling between impulse traffic, neurotransmitter release and synthesis. The evidence concerning the relation between impulse traffic and 5-HT turnover is less clear. TTX (1 μ M) had no effect on 5-HIAA levels in hippocampus in anaesthetised rats⁽²⁹⁾ or caudate-putamen of unanaesthetised rats⁽¹⁷⁾.

Release of transmitter from the nerve terminals on arrival of an action potential depends on the influx of calcium⁽¹⁸⁾. The requirement for calcium has been shown using microdialysis for the release of Ach⁽⁵⁾, DA⁽¹⁴⁾, NA^(20,33), and 5-HT^(2,17). The incomplete abolition of 5-HT and NA release by replacement of calcium with cadmium is again could be due to the fact that not all nerve terminals were reached by the cadmium.

As with TTX, the effect of calcium replacement with cadmium has different

effects on transmitters and metabolites. In the case of DOPAC the reduction is slightly less than that of NA. In noradrenergic neurones calcium in addition to triggering exocytotic NA release, stimulates a pre synaptic adenylate cyclase and is required for activation of tyrosine hydroxylase⁽²⁶⁾ the rate-limiting enzyme for NA synthesis.

By contrast 5-HIAA, which showed little reduction with TTX, is reduced to a greater extent than is 5-HT by cadmium. This suggests an important role for intracellular calcium in 5-HT synthesis. Other work has shown that electrical stimulation of the dorsal raphe increases the level of 5-HT by a mechanism that is attenuated by calcium free perfusate^(28,29), although after replacement of Ca⁺⁺ with Na⁺⁺ ions there is no change in 5-HIAA in caudate putamen⁽¹⁷⁾. Tryptophan hydroxylase, the rate-limiting enzyme in 5-HT synthesis, requires calcium, and an increase in synthesis has an absolute requirement for extracellular calcium ions^(10,11). Thus not only release but also synthesis of both neurotransmitters in hippocampal nerve terminals are calcium dependent.

5-HIAA is the main deaminated metabolite of 5-HT via monoamine oxidase pathway, and represents the turnover of both newly synthesised 5-HT that has not been released and 5-HT which has been taken up by the nerve terminals after release. In some studies 5-HIAA levels have been

used as an index of the activity of serotonergic neurones^(3,24).

Tissue levels of 5-HIAA represent overall 5-HT turnover⁽²⁷⁾; in the presence of a 5-HT re-uptake blocker the re-entry of released 5-HT is prevented and the 5-HIAA levels reflect only the turnover of the newly synthesised 5-HT and not released 5-HT.

5-HIAA levels, in the presence of uptake blocker may thus serve as an index of 5-HT

synthesis. A more direct measurement of the activity of TrH, is the accumulation of 5-HTP in the presence of m-hydroxybenzylhydrazine, the inhibitor for the decarboxylase enzyme⁽⁸⁾. The effects of TTX and cadmium on 5-HIAA in the presence of citalopram suggest that there is basal rate of synthesis, which is largely independent of impulse traffic, but is very sensitive to the extracellular concentration of Ca^{++} .

REFERENCES

1) Aston - Jones, G. and Bloom, F.E. (1981). Norepinephrine - containing locus coeruleus neurons in behaving rats exhibit pronounced responses to non - noxious environmental stimuli - *Journal of Neuroscience* 1: 887.

2) Carboni, E. and di Chiara, G. (1989). Serotonin release estimated by transcortical dialysis in freely-moving rats. *Neuroscience* 32: 637-645.

3) Cespuglio, R., Faradji, H., Ponchon, J.L., Buda, M., Riou, F., Gonon, F., Pujol, J.F. and Jouvot, M. (1981). Differential pulse voltammetry in brain tissue. I. Detection of 5-hydroxyindoles in the rat striatum. *Brain Research* 223: 287-298.

4) Cespuglio, R., Sarda, N., Gharib, A., Faradji, H. and Chastrette, N. (1986). Differential pulse voltammetry in vivo with working carbon fibre electrodes : 5-hydroxyindole compounds or uric acid

detection? *Experimental Brain Research* 64:589-595.

5) Consolo, S.W.C.F., Fiorentini, F., Ladinsky, H. and Vezzani, A. (1987). Determination of endogenous acetylcholine release in freely moving rats by transstriatal dialysis coupled to a radioenzymatic assay: Effect of drugs. *Journal of Neurochemistry* 48: 1459-1465.

6) Damsma, G., Westerink, B.H.C., de Vries, J. B., van Berg, C.J. and Horn, A.S. (1987a). Measurement of acetylcholine release in freely moving rats by means of automated intracerebral dialysis. *Journal of Neurochemistry* 48:1523-1528.

7) Damsma, G., Westerink, B.H.C., Imperato, A., Rollema, H., de Vries, J.B. and Horn, A.S. (1987b). Automated brain dialysis of acetylcholine in freely moving rats: Detection of basal acetylcholine. *Life Sciences* 41: 873-876.

8) De Souza, E.B. and Vanloon, G. (1986). Brain serotonin and catecholamine responses to repeated stress in rats. *Brain Research* 367:77-86.

9) Di Chiara, G.(1990). In vivo brain dialysis of neurotransmitters. *Trends in Pharmacological Sciences* 11: 116-121.

10) Elks, M.L., Youngblood, W.W. and Kizer, J.S. (1979). Synthesis and release of serotonin by brain slices: Effect of ionic manipulations and cationic ionophores. *Brain Research* 172: 461-469.

11) Hamon, M. and Bourgoin, S. (1979). Characterization of the Ca^{2+} -induced proteolytic activation of tryptophan hydroxylase from the rat brain stem. *Journal of Neurochemistry* 32:1837-1844.

12) Hery, F., Simonnet, G., Bourgoin, S., Soubrie, P., Artaud, F., Hamon, M. and Glowinski, J. (1979). Effect of nerve activity on the in vivo release of 3H -Serotonin continuously formed from L - 3H -tryptophan in caudate nucleus of the cat. *Brain Research* 169: 317.

13) Hery, F. and Ternaux, J.P. (1981). Regulation of release processes in central serotonergic neurons. *Journal of Physiology (Paris)* 77:287-301.

14) Imperato, A. and Di Chiara, G. (1984). Trans-striatal dialysis coupled to reverse phase high performance liquid chromatography with electrochemical detection : A new method for the study of the in vivo dopamine release of endogenous

dopamine and metabolites, *Journal of Neuroscience* 4: 966-977.

15) Jacobson, I. and Hamberger, A. (1984). Veratridine-induced release in vivo and in vitro of amino acids in the rabbit olfactory bulb. *Brain Research* 299: 103-112.

16) Kalen, P., Kokaia, M., Lindvall, O. and Bjorklund, A. (1988a). Basic characteristics of noradrenaline release in the hippocampus of intact and 6-hydroxydopamine lesioned rats as studied by in vivo microdialysis. *Brain Research* 474: 374-379.

17) Kalen, P., Strecker, R.E., Rosengren, E. and Bjorklund, A. (1988b). Endogenous release of neuronal serotonin and 5HIAA in the caudate-putamen of rat as revealed by intracerebral dialysis coupled to high performance liquid chromatography with fluorimetric detection. *Journal of Neurochemistry* 51:

18) Katz, B. and Miledi, R. (1965). The effect of calcium on acetylcholine release from motor nerve terminals. *Proceedings of the Royal Society, London B* 161:496-503.

19) Kennett, G. and Joseph, M.H. (1982). Dose in vivo voltammetry in hippocampus measure 5-HT release? *Brain Research* 236:305-316.

20) L'Heureux, R., Dennis, T., Curet, O. and Scatton, B. (1986). Measurement of endogenous noradrenaline release in the rat cerebral cortex in vivo by transcortical

dialysis: Effect of drugs affecting noradrenergic transmission. *Journal of Neurochemistry* 46: 1794-1801.

21) Marsden, C.A., Conti, Y., Strobe, E., Curlzon, G. and Adams, R.N. (1979). Monitoring 5-Hydroxytryptamine release in the brain of the freely moving unanesthetized rat using *in vivo* voltammetry. *Brain Research* 171: 85-99.

22) Marsden, C.A., Martin, K. F., Routledge, C., Brazell, M.P. and Maidment, N.T. (1986). Application of intracerebral dialysis and *in vivo* voltammetry to pharmacological and physiological studies of amine neurotransmitters. *Annals of the New York Academy of Sciences* 473: 106-125.

23) O'Neill, R., Fillenz, M., Grunewald, R.A., Blomfield, M.R., Albery, W.J., Jamieson, C.M., Williams, J.H. and Gray, J.A. (1984). Voltammetric carbon paste electrodes monitor uric acid and not 5HIAA at the 5-hydroxyindole potential in the rat brain. *Neuroscience Letters* 45:39-46.

24) Reinhard, J.F. and Wurtman, R.J. (1977). Relation between brain 5HIAA levels and the release of serotonin in to brain synapses. *Life Sciences* 21: 1741-1746.

25) Reisine, T.D., Soubrie, P., Ferron, A., Blas, C., Romo, R. and Glowinski, J. (1984). Evidence for dopaminergic innervation of the cat lateral habenula: Its role in controlling serotonin transmission in the basal ganglia. *Brain Research* 308: 281-288.

26) Roth, R.H., Morgenroth III, V.H. and

Salzman, P.M. (1975). Tyrosine hydroxylase : Allosteric activation induced by stimulation of central noradrenergic neurons. *Archives of Pharmacology* 289:327-343.

27) Shanon, N.J., Gunnet, J.W. and Moore, K.E. (1986). A comparison of biochemical indices of 5-hydroxytryptaminergic neuronal activity following electrical stimulation of the dorsal raphe nucleus. *Journal of Neurochemistry* 47: 958-965.

28) Sharp, T., Bramwell, S.R., Clark, D. and Graham-Smith, D.G. (1989). *In vivo* measurement of extracellular 5-hydroxytryptamine in hippocampus of the anaesthetized rat using microdialysis: Changes in relation to 5-hydroxytryptaminergic neuronal activity. *Journal of Neurochemistry* 53: 234-240.

29) Sharp, T., Bramwell, S.R. and Graham-Smith, D.G. (1990) Release of endogenous 5-hydroxytryptamine in rat ventral hippocampus evoked by electrical stimulation of the dorsal raphe nucleus as detected by microdialysis: Sensitivity to tetrodotoxin, calcium and calcium antagonists. *Neuroscience* 39: 629-637.

30) Soubrie, P., Reisine, T.D. and Glowinski, J. (1984). Functional aspects of serotonin transmission in the basal ganglia: A review and an *in vivo* approach using push pull cannula technique. *Neuroscience* 13: 605-625.

31) Stamford, J . A . (1985). *In vivo* voltammetry : Promise and perspective. *Brain Research* 10: 119-135.

32) Trulson, M.E. and Jacobs, B.L. (1979). Effects of 5-methoxy-N , N-diethyltryptamine on behavior and raphe unit activity in freely moving cats. *European Journal of Pharmacology* 54: 43-50.

33) Van Veldhuizen, M.J.A., Feentra, M.G.P., Boer, G.J. and Westerink, B.H.C.

(1990). Microdialysis studies on cortical noradrenaline release: Basic characteristic, significance of extracellular calcium and massive post-mortem increase. *Neuroscience Letters* 119:233-236.

34) Westerink, B.H.C., Damsma, G., Rollema, H., de Vries, J.B. and Horn, A.S. (1987). Scope and limitations of *in vivo* brain dialysis: A comparison of its application to various transmitter systems. *Life Sciences* 41: 1763-1776.